

Pediatric Dosing Chart

Dosage for children from your healthcare professional

Use **ONLY** with dosing cup provided with the product. **DO NOT** use any other dosing device.

ASK A DOCTOR before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.


DO NOT USE if you've had an allergic reaction to ZYRTEC® products or ingredients, or to an antihistamine containing hydroxyzine.

Use products **ONLY** as directed. Always read and follow the label on all ZYRTEC® products.

Children's ZYRTEC® Allergy Syrup

Available in: Grape, Bubblegum

AGE	Under 2 yrs	2 to under 6 yrs	6+ yrs
DOSE	Ask a doctor	2.5 mL or 5 mL	5 mL or 10 mL



MAX: 5 mL once daily or 2.5 mL every 12 hours



MAX: 10 mL once daily or 5 mL every 12 hours




Children's ZYRTEC® Chewables 2.5 mg*

Available in: Grape

AGE	Under 2 yrs	2 to 6 yrs
DOSE	Ask a doctor	1 or 2 tablets

MAX: 2 tablets (5 mg) once daily or 1 tablet (2.5 mg) every 12 hours

Chew or crush tablets completely before swallowing.



Children's ZYRTEC® Chewables 10 mg* and Children's ZYRTEC® Dissolve Tabs**

Available in: Grape, Citrus

AGE	Under 6 yrs	6+ yrs
DOSE	Ask a doctor	1 tablet (10 mg) daily

Chewables: Chew or crush tablets completely before swallowing.

Chewables and Dissolve Tabs: A 5 mg product may be appropriate for less severe symptoms.



Children's ZYRTEC® Hives **New**

Available in: Grape

AGE	Under 6 yrs	6+ yrs
DOSE	Ask a doctor	5 mL or 10 mL

MAX: 10 mL once daily or 5 mL every 12 hours



ACTIVE INGREDIENT: Cetirizine

Today's date: _____

This dosing recommendation from your healthcare professional will expire in 14 DAYS.

*Product eligibility may vary by plan. Check with your benefits administrator to determine which expenses are eligible before you shop. **Relieves sneezing, runny nose, itchy/watery eyes and itching of the nose/throat. Use products only as directed. © Kenvue Brands LLC 2025 · ZYR-61037

THE BRAND MOMS TURN TO FIRST*

*When they first treat their children's upper respiratory allergies.



AGE	DOSE
6-11 years	Take 5 mL to 10 mL every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
2-5 years	Do not use unless directed by a doctor.
Under 2 years	Do not use.


AGE	DOSE
12+ years	Take 10 mL every 4 hours. Do not take more than 6 doses in 24 hours.
6-11 years	Take 5 mL every 4 hours. Do not take more than 6 doses in 24 hours.
4-5 years	Do not use unless directed by a doctor.
Under 4 years	Do not use.

AGE	DOSE
12+ years	2 to 4 chewable tablets (25 mg to 50 mg) every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
6-11 years	1 to 2 chewable tablets (12.5 mg to 25 mg) every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
2-5 years	Do not use unless directed by a doctor.
Under 2 years	Do not use.

See other side for dosing information
on Children's ZYRTEC®



Use products only as directed.



Use product only as directed.



Use product only as directed.

Children's BENADRYL®
Allergy Liquid†‡

Children's BENADRYL®
Allergy Plus Congestion†‡

Children's BENADRYL®
Chewables†

For additional information on allergy
management, talk to your pediatrician.

mL = milliliter.

†Relieves runny nose; sneezing; itchy, watery eyes; and itching
of the nose or throat.

‡Use only with dosing cup provided with product.